arXiv:math/0510171v1 [math.GM] 9 Oct 2005

THERE ARE INFINITELY MANY PAIRS OF TWIN PRIME

ZHANLE DU AND SHOUYU DU

ABSTRACT. We proved that there are infinitely many pairs of twin prime.

1. INTRODUCTION

Let P=$\{p_1, p_2, ..., p_v\} = \{2, 3, ..., p_v\}$ be the primes not exceeding $\sqrt{n}$, then the number of primes not exceeding n [1] is,

$$\pi(n) = \left\{ \begin{array}{l} (\pi(\sqrt{n}) - 1) + n - \left(\left\lfloor \frac{n}{p_1} \right\rfloor + \left\lfloor \frac{n}{p_2} \right\rfloor + \cdots + \left\lfloor \frac{n}{p_v} \right\rfloor \right) \\ + \left(\left\lfloor \frac{n}{p_1 p_2} \right\rfloor + \left\lfloor \frac{n}{p_1 p_3} \right\rfloor + \cdots + \left\lfloor \frac{n}{p_{v-1} p_v} \right\rfloor \right) + \cdots . \end{array} \right. \tag{1.1}$$

For simplicity, we can write it as,

$$\begin{array}{rl} \pi(n) & = (\pi(\sqrt{n}) - 1) + n \left[1 - \frac{1}{p_1}\right] \left[1 - \frac{1}{p_2}\right] \cdots \left[1 - \frac{1}{p_v}\right] \\ & = (\pi(\sqrt{n}) - 1) + n \prod_{i=1}^{v} \left[1 - \frac{1}{p_i}\right], \end{array} \tag{1.2}$$

where $n\left[1 - \frac{1}{p_i}\right] = n - \left\lfloor \frac{n}{p_i} \right\rfloor$, $n\left[1 - \frac{1}{p_i}\right]\left[1 - \frac{1}{p_j}\right] = n - \left\lfloor \frac{n}{p_i} \right\rfloor - \left\lfloor \frac{n}{p_j} \right\rfloor + \left\lfloor \frac{n}{p_i p_j} \right\rfloor$. The operator $\left[1 - \frac{1}{p_i}\right]$ will leave the items which are not multiples of p_i.

In this paper, $\lfloor x \rfloor \le x$ is the floor function, $\lceil x \rceil \ge x$ the ceiling function of x. The integral operator '[ ]', which is not the floor function in this paper, has meanings only operating on (real) number: $m[\,] = [m] = \lfloor m \rfloor$.

If p_t and $p_t + 2$ are both primes, then this prime pair is called twin primes. Is there infinitely twin primes? It is a long-term unsolved conjecture[1, 2, 3, 4]. We will proved it as theorem 1.1.

Theorem 1.1. *There are infinitely many pairs of twin prime.*

2. THE NUMBER OF TWIN PRIMES FROM p_v^2 TO p_{v+1}^2

Let $Z = \{1, 2, \cdots, n\} (n < p_{v+1}^2)$ be a natural arithmetic progression, $Z' = Z + 2 = \{3, 5, \cdots, n + 2\}$ be its accompanying arithmetic progression, so that $Z'_k = Z_k + 2, k = 1, 2, \cdots, n$. There are n such pairs.

$$\left\{ \begin{array}{lllllll} Z & = \{ & 1, & 2, & \cdots, & n & \} \\ Z' & = \{ & 3, & 5, & \cdots, & n+2 & \}. \end{array} \right. \tag{2.1}$$

Date: Oct.3, 2005.

2000 *Mathematics Subject Classification.* 11A41.

Key words and phrases. prime,twin primes, floor function, ceiling function, integral operator.

All rights reserved. This is one of a serial works, and was supported financially by Pro. Wei WANG and on loan in part.

Corresponding author: Zhanle Du.

If we delete all the pairs in which one or both items Z_k and Z'_k are composite integers of all primes $p_i \leq \sqrt{n}$, then the pairs left are all twin primes.

For a given p_i, first we delete the multiples of p_i in set Z, or the items of $Z_k \mathbf{mod} p_i = 0$,

$$y(p_i) = \left\lfloor \frac{n}{p_i} \right\rfloor \equiv n \left[\frac{1}{p_i} \right]. \tag{2.2}$$

Secondly we delete the multiples of p_i in Z', i.e. the items of $Z'_k \mathbf{mod} p_i = 0$, or $Z_k \mathbf{mod} p_i = p_i - 2$,

$$y'(p_i) = \left\lfloor \frac{n+2}{p_i} \right\rfloor \equiv n \left[\frac{\widetilde{1}}{p_i} \right]. \tag{2.3}$$

Because $Z_k \mathbf{mod} p_i \neq Z'_k \mathbf{mod} p_i$ when $p_i \neq 2$, the items deleted by $y(p_i)$ and $y'(p_i)$ are not the same. For $p_i = 2$, $Z_k \mathbf{mod} p_i = Z'_k \mathbf{mod} p_i$, we need only delete the items in set Z:

$$y'(p_i) = 0, \quad \text{for} \quad p_i = 2. \tag{2.4}$$

Eq. (2.2) will delete all pairs with $Z_k \mathbf{mod} p_i = 0$ in set Z, and Eq. (2.3) will delete all pairs with $Z_k \mathbf{mod} p_i = p_i - 2$ in set Z.

$$y'(p_i) = n \left[\frac{\widetilde{1}}{p_i} \right] = n \left\lfloor \frac{1}{p_i} \right\rfloor + \delta, \tag{2.5}$$

where $0 \leq \delta \leq 1$ with,

$$\delta = \begin{cases} 0: & 0 \leq n \mathbf{mod} p_i + 2 < p_i \\ 1: & \text{else.} \end{cases} \tag{2.6}$$

The pairs left, when deleted all the multiples of p_i in both Z and Z', have,

$$M(p_i) = n - y(p_i) - y'(p_i) \equiv n \left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i} \right]. \tag{2.7}$$

The operator $\left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right]$, when operating on n, will leaves the items having not those of $Z_k \mathbf{mod} p_i = 0, p_i - 2$.

For another prime p_j, we should also delete its multiples in both Z and Z', but if there is an item which is composite integer of both p_i and p_j then this item should be deleted only once. Thus $n \left[\frac{1}{p_j}\right] \left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right]$ will delete the multiples of p_j in Z, which have not been deleted by p_i. $n \left[\frac{\widetilde{1}}{p_j}\right] \left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right]$ will delete the multiples of p_j in Z', which have not been deleted by p_i. The pairs will leave,

$$\begin{aligned} M(p_i, p_j) &= n \left[1 - \tfrac{1}{p_i} - \tfrac{\widetilde{1}}{p_i}\right] - n \left[\tfrac{1}{p_j}\right] \left[1 - \tfrac{1}{p_i} - \tfrac{\widetilde{1}}{p_i}\right] - n \left[\tfrac{\widetilde{1}}{p_j}\right] \left[1 - \tfrac{1}{p_i} - \tfrac{\widetilde{1}}{p_i}\right] \\ &\equiv n \left[1 - \tfrac{1}{p_i} - \tfrac{\widetilde{1}}{p_i}\right] \left[1 - \tfrac{1}{p_j} - \tfrac{\widetilde{1}}{p_j}\right] \\ &= n \left[1 - \tfrac{1}{p_i} - \tfrac{\widetilde{1}}{p_i} - \tfrac{1}{p_j} + \tfrac{1}{p_i p_j} + \tfrac{\widetilde{1}}{p_i} \tfrac{1}{p_j} - \tfrac{\widetilde{1}}{p_j} + \tfrac{1}{p_i} \tfrac{\widetilde{1}}{p_j} + \tfrac{\widetilde{1}}{p_i} \tfrac{\widetilde{1}}{p_j}\right], \end{aligned} \tag{2.8}$$

the meaning is as follows,

$$
\left\{
\begin{array}{l}
n\left[\frac{1}{p_i}\right] = \left\lfloor \frac{n}{p_i} \right\rfloor \\
n\left[\frac{\widetilde{1}}{p_i}\right] = \left\lfloor \frac{n+2}{p_i} \right\rfloor \\
n\left[\frac{1}{p_i}\right]\left[\frac{1}{p_j}\right] = n\left[\frac{1}{p_ip_j}\right] = \left\lfloor \frac{n}{p_ip_j} \right\rfloor \\
n\left[\frac{\widetilde{1}}{p_i}\right]\left[\frac{\widetilde{1}}{p_j}\right] = n\left[\frac{\widetilde{1}}{p_ip_j}\right] = \left\lfloor \frac{n+2}{p_ip_j} \right\rfloor \\
n\left[\frac{1}{p_i}\right]\left[\frac{\widetilde{1}}{p_j}\right] = n\left[\frac{1}{p_i}\frac{\widetilde{1}}{p_j}\right] = \left\lfloor \frac{n+\theta_{i,j}}{p_ip_j} \right\rfloor = \left\lfloor \frac{n+p_ip_j-\lambda_{i,j}}{p_ip_j} \right\rfloor,
\end{array}
\right.
\tag{2.9}
$$

where $1 \leq \lambda_{i,j} \leq p_ip_j$ is the first (number) position in Z with $p_i|Z_{\lambda_{i,j}}$ and $p_j|Z'_{\lambda_{i,j}}$,

$$
\left\{
\begin{array}{l}
Z_{\lambda_{i,j}}\mathbf{mod}p_i = 0 \\
Z_{\lambda_{i,j}}\mathbf{mod}p_j = p_j - 2.
\end{array}
\right.
\tag{2.10}
$$

Let $X = \{\lambda p_i, \lambda = 1, 2, \cdots, \left\lfloor \frac{n}{p_i} \right\rfloor\}$, λ_j be the first item with $\lambda_j p_i\mathbf{mod}p_j = p_j - 2$, then $\lambda_{i,j} = \lambda_j p_i$,

$$
n\left[\frac{1}{p_i}\right]\left[\frac{\widetilde{1}}{p_j}\right] = \left\lfloor \frac{n+p_ip_j-\lambda_{i,j}}{p_ip_j} \right\rfloor = \left\lfloor \frac{\left\lfloor \frac{n}{p_i} \right\rfloor + p_j - \lambda_j}{p_j} \right\rfloor.
\tag{2.11}
$$

If there is no such $\lambda_{i,j}$ in Z, i.e., $\lambda_{i,j} \geq (n+1)$, then the last item in Eq. (2.9) will equal zero. Because $1 \leq \lambda_j \leq (p_j - 1)$, so $p_i \leq \theta_{i,j} = p_ip_j - \lambda_{i,j} \leq p_i(p_j - 1)$.

When the multiples of all primes $p_i \leq p_v < \sqrt{n}$ in both Z and Z' have been deleted, the pairs left will be prime pairs and have,

$$
\begin{aligned}
D_0(n) &= n\left[1 - \frac{1}{p_1}\right]\left[1 - \frac{1}{p_2} - \frac{\widetilde{1}}{p_2}\right]\cdots\left[1 - \frac{1}{p_v} - \frac{\widetilde{1}}{p_v}\right] \\
&= n\left[1 - \frac{1}{p_1}\right]\prod_{i=2}^{v}\left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right],
\end{aligned}
\tag{2.12}
$$

The total number of twin primes in n is,

$$
\begin{aligned}
D(n) &= D_0(n) + D(\sqrt{n}) - D_1 \\
&= n\left[1 - \frac{1}{p_1}\right]\prod_{i=2}^{v}\left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right] + D(\sqrt{n}) - D_1,
\end{aligned}
\tag{2.13}
$$

where $D(\sqrt{n}) \geq 0$ is the number of twin primes $[p_i, p_{i+1}]$ when $p_i \leq p_v < \sqrt{n}$, where $p_{i+1} = p_i + 2$ and,

$$
D_1 = \left\{
\begin{array}{ll}
1 & \text{if } n + 2 = p_{v+1}^2 \text{ and } n = prime \\
0 & \text{else.}
\end{array}
\right.
\tag{2.14}
$$

Because $n < p_{v+1}^2$, if $n = p_{v+1}^2 - 1$ then $n\mathbf{mod}2 = 0$; if $n = p_{v+1}^2 - 2$ and n is prime then $[n, n+2]$ is not twin prime.

Besides, the pair [1,3] has been already deleted since $1\mathbf{mod}p_2 = p_2 - 2$ for $p_2 = 3$.

Example 2.1. $n = 41$, $Z = [1, 2, \cdots, 41]$, $p_i = [2, 3, 5]$,

$\overrightarrow{Z_k\mathbf{mod}2 \neq 0}$ $[1, 3, \cdots, 41]$

$\overrightarrow{Z_k\mathbf{mod}3 \neq 0}$ $[1, 5, 7, 11, 13, 17, 19, 23, 25, 29, 31, 35, 37, 41]$

$\overrightarrow{Z_k\mathbf{mod}5 \neq 0}$ $[1, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41]$

$\overrightarrow{Z_k\mathbf{mod}3 \neq 1}$ $[11, 17, 23, 29, 41]$

$\overrightarrow{Z_k\mathbf{mod}5 \neq 3}$ $[11, 17, 29, 41]$.

So $D_0(41) = 4$.

We can check it from Eq. (2.12) directly.

$$\begin{aligned}
D_0(41) &= n\left[1-\tfrac{1}{2}\right]\left[1-\tfrac{1}{3}-\tfrac{\widetilde{1}}{3}\right]\left[1-\tfrac{1}{5}-\tfrac{\widetilde{1}}{5}\right]\\
&= 41\Big[1-\tfrac{1}{2}-\tfrac{1}{3}+\tfrac{1}{6}-\tfrac{\widetilde{1}}{3}+\tfrac{1}{2}\tfrac{\widetilde{1}}{3}-\tfrac{1}{5}+\tfrac{1}{10}+\tfrac{1}{15}-\tfrac{1}{30}+\tfrac{1}{5}\tfrac{\widetilde{1}}{3}-\tfrac{1}{10}\tfrac{\widetilde{1}}{3}\\
&\quad -\tfrac{\widetilde{1}}{5}+\tfrac{1}{2}\tfrac{\widetilde{1}}{5}+\tfrac{1}{3}\tfrac{\widetilde{1}}{5}-\tfrac{1}{6}\tfrac{\widetilde{1}}{5}+\tfrac{\widetilde{1}}{15}-\tfrac{1}{2}\tfrac{\widetilde{1}}{15}\Big]\\
&= 41-\left\lfloor\tfrac{41}{2}\right\rfloor-\left\lfloor\tfrac{41}{3}\right\rfloor+\left\lfloor\tfrac{41}{6}\right\rfloor-\left[\tfrac{\widetilde{41}}{3}\right]+\left[\tfrac{41}{2}\tfrac{\widetilde{1}}{3}\right]\\
&\quad -\left\lfloor\tfrac{41}{5}\right\rfloor+\left\lfloor\tfrac{41}{10}\right\rfloor+\left\lfloor\tfrac{41}{15}\right\rfloor-\left\lfloor\tfrac{41}{30}\right\rfloor+\left[\tfrac{41}{5}\tfrac{\widetilde{1}}{3}\right]-\left[\tfrac{41}{10}\tfrac{\widetilde{1}}{3}\right]\\
&\quad -\left[\tfrac{\widetilde{41}}{5}\right]+\left[\tfrac{41}{2}\tfrac{\widetilde{1}}{5}\right]+\left[\tfrac{41}{3}\tfrac{\widetilde{1}}{5}\right]-\left[\tfrac{41}{6}\tfrac{\widetilde{1}}{5}\right]+\left[\tfrac{\widetilde{41}}{15}\right]-\left[\tfrac{41}{2}\tfrac{\widetilde{1}}{15}\right]\\
&= 41-20-13+6-\left[\tfrac{41+2}{3}\right]+\left\lfloor\frac{\lfloor\frac{41}{2}\rfloor+3-2}{3}\right\rfloor\\
&\quad -8+4+2-1+\left\lfloor\frac{\lfloor\frac{41}{5}\rfloor+3-2}{3}\right\rfloor-\left\lfloor\frac{\lfloor\frac{41}{10}\rfloor+3-1}{3}\right\rfloor\\
&\quad -\left[\tfrac{41+2}{5}\right]+\left\lfloor\frac{\lfloor\frac{41}{2}\rfloor+5-3}{5}\right\rfloor+\left\lfloor\frac{\lfloor\frac{41}{3}\rfloor+5-1}{5}\right\rfloor-\left\lfloor\frac{\lfloor\frac{41}{6}\rfloor+5-3}{5}\right\rfloor+\left[\tfrac{41+2}{15}\right]-\left\lfloor\frac{\lfloor\frac{41}{2}\rfloor+15-14}{15}\right\rfloor\\
&= 14-14+\left\lfloor\tfrac{21}{3}\right\rfloor\\
&\quad -3+\left\lfloor\tfrac{9}{3}\right\rfloor-\left\lfloor\tfrac{6}{3}\right\rfloor\\
&\quad -8+\left\lfloor\tfrac{22}{5}\right\rfloor+\left\lfloor\tfrac{17}{5}\right\rfloor-\left\lfloor\tfrac{8}{5}\right\rfloor+\left[\tfrac{41+2}{15}\right]-\left\lfloor\tfrac{21}{15}\right\rfloor\\
&= 7+(-3+3-2)+(-8+4+3-1+2-1)\\
&= 4.
\end{aligned}$$

It is the same as before. From $D(\sqrt{41})=2$ for $p_i\le 5$: (3,5) and (5,7), $D_1(n)=0(n+2=43<p_4^2=49)$. So $D(41)=D_0(41)+D(\sqrt{41})-D_1=4+2-0=6$.

The set: $Z=\{3,5,11,17,29,41\}, Z'=Z+2=\{5,7,13,19,31,43\}$. The six twin primes are $(3,5),(5,7),(11,13),(17,19),(29,31),(41,43)$. □

Definition 2.2. The items of $Z_k\mathbf{mod}p_i=0$, or the multiples of p_i, have,

$$S(m,p_i\parallel 0):=\sum_{Z_k\mathbf{mod}p_i=0}1=m\left[\frac{1}{p_i}\right]=\left\lfloor\frac{m}{p_i}\right\rfloor. \tag{2.15}$$

So the items of $Z_k\mathbf{mod}p_j\neq 0,p_j-2$ have,

$$S(m,p_j\nparallel 0,p_j-2)\quad:=\sum_{Z_k\mathbf{mod}p_i\neq 0,p_j-2}1=m\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]. \tag{2.16}$$

Definition 2.3. Let $X=\{X_1,X_2,\cdots,X_t\}$ be an (any) integer set, then

$$t\left[1-\frac{1'}{p_i}-\frac{1''}{p_i}\right]:=\sum_{X_k\mathbf{mod}p_i\neq 0,p_i-2}1=t-\left[\frac{t+\phi_i'}{p_i}\right]-\left[\frac{t+\phi_i''}{p_i}\right]\ge 0 \tag{2.17}$$

is the number of items left after deleted the items of $X_k\mathbf{mod}p_i=0,p_i-2$.

Because $m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]\neq\left(m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\right)\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]$, We can express it as

$$\begin{aligned}
S(m,p_i\nparallel 0,p_i-2;p_j\nparallel 0,p_j-2) &= m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]\\
&= \left(m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\right)\left[1-\frac{1'}{p_j}-\frac{1''}{p_j}\right],
\end{aligned} \tag{2.18}$$

is the number of items left when we first delete those $Z_k\mathbf{mod}p_i=0,p_i-2$ from set Z, and then delete those $X_{k'}\mathbf{mod}p_j=0,p_j-2$ from set $X=\{Z,X_{k'}\mathbf{mod}p_i\neq$

$0, p_i-2, k'=1,2,\cdots, m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\}$, where set X is no longer an arithmetic sequence. In general,

$$m\prod_{i=1}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]=\left(m\prod_{i=1}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\right)\left[1-\frac{1'}{p_j}-\frac{1''}{p_j}\right]. \tag{2.19}$$

3. Some property

$$\begin{cases}\left[1-\frac{1}{p_i}\right]\left[1-\frac{1}{p_j}\right]=\left[1-\frac{1}{p_j}\right]\left[1-\frac{1}{p_i}\right]\\ \left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]=\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right].\end{cases} \tag{3.1}$$

$$\left[\frac{m}{p_i}\right]=\left[\frac{m_1+m_2}{p_i}\right]=\left[\frac{m_1}{p_i}\right]+\left[\frac{m_2}{p_i}\right]+\left[\frac{m_1\mathbf{mod}p_i+m_2\mathbf{mod}p_i}{p_i}\right]. \tag{3.2}$$

$$m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]=ap_i\left(1-\frac{2}{p_i}\right)+b\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\quad \text{for } m=ap_i+b. \tag{3.3}$$

$$-2\le(m_1+m_2)\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]-m_1\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]-m_2\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\le 1. \tag{3.4}$$

Proof. Let $\alpha=m_1\mathbf{mod}p_i, \beta=m_2\mathbf{mod}p_i$,

$$\begin{aligned}&\delta_{12}=(m_1+m_2)\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]-m_1\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]-m_2\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\\ &=m_1+m_2-\left[\frac{m_1+m_2}{p_i}\right]-\left[\frac{m_1+m_2+2}{p_i}\right]-m_1+\left[\frac{m_1}{p_i}\right]+\left[\frac{m_1+2}{p_i}\right]\\ &\quad -m_2+\left[\frac{m_2}{p_i}\right]+\left[\frac{m_2+2}{p_i}\right]\\ &=-\left[\frac{\alpha+\beta}{p_i}\right]-\left[\frac{\alpha+\beta+2}{p_i}\right]+\left[\frac{\alpha+2}{p_i}\right]+\left[\frac{\beta+2}{p_i}\right].\end{aligned}$$

The minimum: $min(\delta_{12})=-1-1=-2$, when $\alpha+\beta\ge p_i$, $\alpha+2<p_i$, $\beta+2<p_i$, and $\alpha+\beta+2\ge p_i$.

The maximum: $max(\delta_{12})=-0-1+1+1=1$, when $\alpha+\beta<p_i$, $\alpha+2\ge p_i$, $\beta+2\ge p_i$. □

We can represent it as

$$\begin{aligned}&(m_1+m_2)\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]=m_1+m_2-\left[\frac{m_1+m_2}{p_i}\right]-\left[\frac{m_1+m_2+2}{p_i}\right]\\ =\ &m_1-\left[\frac{m_1}{p_i}\right]-\left[\frac{m_1+2}{p_i}\right]+m_2-\left[\frac{m_1\mathbf{mod}p_i+m_2}{p_i}\right]-\left[\frac{m_2+(m_1+2)\mathbf{mod}p_i}{p_i}\right]\\ =\ &m_1\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]+m_2\left[1-\frac{1'}{p_i}-\frac{1''}{p_i}\right].\end{aligned} \tag{3.5}$$

$m_2\left[\frac{1'}{p_i}\right]$ will delete the items of $X_k\mathbf{mod}p_i=0$ in the sequence of $X=\{m_1+1, m_1+2,\cdots,m_1+m_2\}$, and $m_2\left[\frac{1''}{p_i}\right]$ will delete the items of $X'_k\mathbf{mod}p_i=0$ in the sequence of $X'=\{X+2\}$ or the items of $X_k\mathbf{mod}p_i=p_i-2=p_i-2$ in set X.

For any m_2, $0\le m_2\left[1-\frac{1'}{p_i}-\frac{1''}{p_i}\right]\le m_2$, so,

$$m'\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\ge m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\quad \text{for } m'\ge m. \tag{3.6}$$

4. Some lemma

Lemma 4.1. *For $m \geq p_i$,*

$$m\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \geq \left\lceil m\left(1-\tfrac{3}{p_i}\right)\right\rceil. \tag{4.1}$$

Proof.

$$\begin{aligned} left &= \left\lceil m-\left\lfloor\tfrac{m}{p_i}\right\rfloor-\left\lfloor\tfrac{m+\theta_i}{p_i}\right\rfloor\right\rceil \geq \left\lceil m-\left\lfloor\tfrac{m}{p_i}\right\rfloor-\left\lfloor\tfrac{m}{p_i}\right\rfloor-1\right\rceil \\ &\geq \left\lceil m-\tfrac{2m}{p_i}-\tfrac{m}{p_i}\right\rceil = right. \end{aligned}$$

□

Lemma 4.2.

$$\begin{cases} -1 \leq & m\left[\tfrac{1}{p_j}\right]\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]-\left\lfloor\tfrac{m}{p_j}\right\rfloor\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \quad \leq 1 \\ -1 \leq & m\left[\tfrac{\widetilde{1}}{p_j}\right]\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]-\left\lceil\tfrac{m+2}{p_j}\right\rceil\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \quad \leq 1. \end{cases} \tag{4.2}$$

Proof.

$$\begin{aligned} \varepsilon_1 &= m\left[\tfrac{1}{p_j}\right]\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]-\left\lfloor\tfrac{m}{p_j}\right\rfloor\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \\ &= \left\lfloor\tfrac{m}{p_j}\right\rfloor-\left\lfloor\tfrac{m}{p_ip_j}\right\rfloor-\left\lfloor\tfrac{m+\theta_{j,i}}{p_ip_j}\right\rfloor-\left\lfloor\tfrac{m}{p_j}\right\rfloor+\left\lfloor\tfrac{m}{p_ip_j}\right\rfloor+\left\lfloor\frac{\left\lfloor\frac{m}{p_j}\right\rfloor+2}{p_i}\right\rfloor \\ &= -\left\lfloor\tfrac{m+\theta_{j,i}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{m+2p_j}{p_ip_j}\right\rfloor \\ &= -\left\lfloor\tfrac{m\,\mathbf{mod}\,p_ip_j+\theta_{j,i}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{m\,\mathbf{mod}\,p_ip_j+2p_j}{p_ip_j}\right\rfloor, \end{aligned}$$

so that $-1 \leq \varepsilon_1 \leq 1$.

$$\begin{aligned} \varepsilon_2 &= m\left[\tfrac{\widetilde{1}}{p_j}\right]\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]-\left\lfloor\tfrac{m+2}{p_j}\right\rfloor\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \\ &= \left\lfloor\tfrac{m+2}{p_j}\right\rfloor-\left\lfloor\tfrac{m+\theta_{i,j}}{p_ip_j}\right\rfloor-\left\lfloor\tfrac{m+2}{p_ip_j}\right\rfloor-\left\lfloor\tfrac{m+2}{p_j}\right\rfloor+\left\lfloor\left\lfloor\tfrac{m+2}{p_j}\right\rfloor\tfrac{1}{p_i}\right\rfloor+\left\lfloor\frac{\left\lfloor\frac{m+2}{p_j}\right\rfloor+2}{p_i}\right\rfloor \\ &= -\left\lfloor\tfrac{m\,\mathbf{mod}\,p_ip_j+\theta_{i,j}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{m\,\mathbf{mod}\,p_ip_j+2+2p_j}{p_ip_j}\right\rfloor, \end{aligned}$$

so that $-1 \leq \varepsilon_2 \leq 1$. □

Lemma 4.3. *For $m \geq p_j^2$, $p_j > p_i \geq 3$,*

$$m\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right] \geq \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]. \tag{4.3}$$

$$m\left[1-\tfrac{1}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right] \geq \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\left[1-\tfrac{1}{p_i}\right] \quad \textit{for } p_i = 2. \tag{4.4}$$

Proof of (4.3). Let $m = sp_ip_j + t$, $t = ap_j + b$, $0 \leq a \leq (p_i - 1)$, $0 \leq b \leq (p_j - 1)$, Because $m \geq p_j^2$, $p_j > p_i$, so $s \geq 1$. From Eq. (3.2), (3.3),

$$\begin{aligned}
\varepsilon \quad &= m\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]-\left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\\
&= sp_ip_j\left(1-\tfrac{2}{p_i}\right)\left(1-\tfrac{2}{p_j}\right)-sp_ip_j\left(1-\tfrac{2}{p_i}\right)\left(1-\tfrac{3}{p_j}\right)\\
&\quad +t-\left\lfloor\tfrac{t}{p_i}\right\rfloor-\left\lfloor\tfrac{t+2}{p_i}\right\rfloor-\left\lfloor\tfrac{t}{p_j}\right\rfloor+\left\lfloor\tfrac{t}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{t+\theta_{j,i}}{p_ip_j}\right\rfloor-\left\lfloor\tfrac{t+2}{p_j}\right\rfloor+\left\lfloor\tfrac{t+\theta_{i,j}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{t+2}{p_ip_j}\right\rfloor\\
&\quad -\left(t-\left\lfloor\tfrac{3t}{p_j}\right\rfloor\right)+\left\lfloor\frac{t-\left\lfloor\frac{3t}{p_j}\right\rfloor}{p_i}\right\rfloor+\left\lfloor\frac{t+2-\left\lfloor\frac{3t}{p_j}\right\rfloor}{p_i}\right\rfloor\\
&= s(p_i-2)\\
&\quad +\left\lfloor\tfrac{3t}{p_j}\right\rfloor-\left\lfloor\tfrac{t}{p_j}\right\rfloor-\left\lfloor\tfrac{t+2}{p_j}\right\rfloor+\left\lfloor\tfrac{t}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{t+\theta_{j,i}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{t+\theta_{i,j}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{t+2}{p_ip_j}\right\rfloor\\
&\quad -\left\lfloor\frac{\left\lfloor\frac{3t}{p_j}\right\rfloor}{p_i}\right\rfloor-\varepsilon_1-\left\lfloor\frac{\left\lfloor\frac{3t}{p_j}\right\rfloor}{p_i}\right\rfloor-\varepsilon_2\\
&= s(p_i-2)\\
&\quad +3a+\left\lfloor\tfrac{3b}{p_j}\right\rfloor-a-a-\left\lfloor\tfrac{b+2}{p_j}\right\rfloor+0+\left\lfloor\tfrac{ap_j+b+\theta_{j,i}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{ap_j+b+\theta_{i,j}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{ap_j+b+2}{p_ip_j}\right\rfloor\\
&\quad -2\left\lfloor\frac{3a+\left\lfloor\frac{3b}{p_j}\right\rfloor}{p_i}\right\rfloor-\varepsilon_1-\varepsilon_2\\
&= s(p_i-2)+a+\left\lfloor\tfrac{3b}{p_j}\right\rfloor+\left\lfloor\tfrac{ap_j+b+\theta_{j,i}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{ap_j+b+\theta_{i,j}}{p_ip_j}\right\rfloor+\left\lfloor\tfrac{ap_j+b+2}{p_ip_j}\right\rfloor\\
&\quad -\varepsilon_1-\varepsilon_2-2\varepsilon_3-\varepsilon_4,
\end{aligned}$$

where

$$\begin{cases}
\varepsilon_1 &= \left\lfloor\frac{\left\lfloor\frac{3t}{p_j}\right\rfloor\mathbf{mod}p_i+\left(t-\left\lfloor\frac{3t}{p_j}\right\rfloor\right)\mathbf{mod}p_i}{p_i}\right\rfloor\le 1\\
\varepsilon_2 &= \left\lfloor\frac{\left\lfloor\frac{3t}{p_j}\right\rfloor\mathbf{mod}p_i+\left(t+2-\left\lfloor\frac{3t}{p_j}\right\rfloor\right)\mathbf{mod}p_i}{p_i}\right\rfloor\le 1\\
\varepsilon_3 &= \left\lfloor\frac{3a+\left\lfloor\frac{3b}{p_j}\right\rfloor}{p_i}\right\rfloor\le 2\\
\varepsilon_4 &= \left\lfloor\frac{b+2}{p_j}\right\rfloor\le 1,
\end{cases}$$

$$\Delta\varepsilon=\varepsilon_1+\varepsilon_2+2\varepsilon_3+\varepsilon_4\le 7.$$

(1) If $p_i\ge 11$, then $\varepsilon\ge s(11-2)-\Delta\varepsilon\ge 9-7>0$.
(2) If $p_i=7$, then
  (a) If $a\ge 2$, then $\varepsilon\ge s(p_i-2)+a-\Delta\varepsilon\ge 5+2-7=0$.
  (b) If $a\le 1$, then $\varepsilon_3=\left\lfloor\frac{3a+\left\lfloor\frac{3b}{p_j}\right\rfloor}{p_i}\right\rfloor\le\left\lfloor\frac{3+2}{7}\right\rfloor=0$, $\Delta\varepsilon\le 3$, so $\varepsilon\ge s(p_i-2)-\Delta\varepsilon\ge 5-3>0$.
(3) If $p_i=5$, then
  (a) If $a\ge 4$, then $\varepsilon\ge s(p_i-2)+a-\Delta\varepsilon\ge 3+4-7=0$.
  (b) If $a=0$, then $\varepsilon_3=0$, $\Delta\varepsilon\le 3$, so $\varepsilon\ge s(p_i-2)-\Delta\varepsilon\ge 3-3=0$.
  (c) If $a=1$: if $\left\lfloor\frac{3b}{p_j}\right]\ge 1$ then $\varepsilon_3\le 1$, $\Delta\varepsilon\le 5$, so $\varepsilon\ge s(p_i-2)+a+\left\lfloor\frac{3b}{p_j}\right]-\Delta\varepsilon\ge 3+1+1-5=0$; else for $\left\lfloor\frac{3b}{p_j}\right]=0$ then $\varepsilon_3=0$, $\Delta\varepsilon\le 3$, so $\varepsilon\ge s(p_i-2)+a-\Delta\varepsilon\ge 3+1-3>0$.
  (d) If $a=2$, then $\varepsilon_3=1$, $\Delta\varepsilon\le 5$, so $\varepsilon\ge s(p_i-2)+a-\Delta\varepsilon\ge 3+2-5=0$.

(e) If $a = 3$: if $\left\lfloor \frac{3b}{p_j} \right\rfloor \geq 1$ then $\varepsilon \geq s(p_i - 2) + a + \left\lfloor \frac{3b}{p_j} \right\rfloor - \Delta\varepsilon \geq 3 + 3 + 1 - 7 = 0$; else for $\left\lfloor \frac{3b}{p_j} \right\rfloor = 0$ then $\varepsilon_3 = \left\lfloor \frac{3a + \left\lfloor \frac{3b}{p_j} \right\rfloor}{p_i} \right\rfloor = 1$, $\Delta\varepsilon \leq 5$, so $\varepsilon \geq s(p_i - 2) + a - \Delta\varepsilon \geq 3 + 3 - 5 > 0$.

(4) If $p_i = 3$, then

$$\begin{cases} a - 2\varepsilon_3 & = a - 2\left\lfloor \frac{3a + \left\lfloor \frac{3b}{p_j} \right\rfloor}{3} \right\rfloor = a - 2a = -a \\ \varepsilon_1 & = \left\lfloor \frac{\left(3a + \left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3 + \left(ap_j + b - 3a - \left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3}{3} \right\rfloor \\ & = \left\lfloor \frac{\left(\left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3 + \left(ap_j + b - \left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3}{3} \right\rfloor \\ \varepsilon_2 & = \left\lfloor \frac{\left(3a + \left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3 + \left(ap_j + b + 2 - 3a - \left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3}{3} \right\rfloor \\ & = \left\lfloor \frac{\left(\left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3 + \left(ap_j + b + 2 - \left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3}{3} \right\rfloor, \end{cases}$$

$$\begin{aligned} \varepsilon &= s - a + \left\lfloor \frac{3b}{p_j} \right\rfloor + \left\lfloor \frac{ap_j + b + \theta_{j,i}}{p_i p_j} \right\rfloor + \left\lfloor \frac{ap_j + b + \theta_{i,j}}{p_i p_j} \right\rfloor + \left\lfloor \frac{ap_j + b + 2}{p_i p_j} \right\rfloor \\ &\quad - \varepsilon_1 - \varepsilon_2 - \varepsilon_4. \end{aligned}$$

(a) If $\left\lfloor \frac{3b}{p_j} \right\rfloor = 0$, then $\varepsilon_1 = \varepsilon_2 = 0$, $b < p_j/3$, $b + 2 < p_j/3 + 2 = (p_j + 6)/3 < p_j$, so $\varepsilon_4 = 0$. If $a \leq 1$ then $\varepsilon \geq s - a \geq 0$. Else for $a = 2 (< p_i)$, because $\theta_{j,i} = p_i p_j - \lambda p_j \geq p_j(1 \leq \lambda \leq p_i - 1)$, $\left\lfloor \frac{ap_j + b + \theta_{j,i}}{p_i p_j} \right\rfloor \geq \left\lfloor \frac{2p_j + p_j}{3p_j} \right\rfloor = 1$. So $\varepsilon \geq s - a + 1 \geq 0$.

(b) If $\left\lfloor \frac{3b}{p_j} \right\rfloor \geq 1$, then $\varepsilon' = s + \left\lfloor \frac{ap_j + b + \theta_{i,j}}{3p_j} \right\rfloor - \varepsilon_4 \geq 1$.

Proof.
- If $s \geq 2$ then $\varepsilon' \geq 1$.
- If $s = 1$, i.e., $s = \left\lfloor \frac{m}{p_i p_j} \right\rfloor \geq \left\lfloor \frac{p_j^2}{3p_j} \right\rfloor = 1$, so $p_j = 5$. From $5^2 = 3 \cdot 5 + 2 \cdot 5$, we have $a = 2$. Let us consider $\theta_{i,j} = p_i p_j - \lambda_{i,j}$, $\lambda_{i,j} = \lambda p_i = 3\lambda$, with the condition $(3\lambda + 2)\mathbf{mod}5 = 0$ i.e. $\lambda = 1$, $\theta_{i,j} = p_i p_j - 3\lambda = 15 - 3 = 12$. $\left\lfloor \frac{ap_j + b + \theta_{i,j}}{p_i p_j} \right\rfloor \geq \left\lfloor \frac{2 \cdot 5 + 12}{3 \cdot 5} \right\rfloor = 1$, Therefore,

$$\varepsilon' = s + \left\lfloor \frac{ap_j + b + \theta_{i,j}}{3p_j} \right\rfloor - \varepsilon_4 \geq 1 \quad \text{if} \quad s = 1.$$

□

Besides, $\varepsilon'' = \left\lfloor \frac{3b}{p_j} \right\rfloor - \varepsilon_1 - \varepsilon_2 \geq 0$.

Proof.
- if $\left\lfloor \frac{3b}{p_j} \right\rfloor = 2$, then $\varepsilon'' \geq 0$.
- if $\left\lfloor \frac{3b}{p_j} \right\rfloor = 1$, because,

$$\begin{aligned}\varepsilon_1+\varepsilon_2 &= \left\lfloor \tfrac{1+(ap_j+b-1)\mathbf{mod}3}{3}\right\rfloor + \left\lfloor \tfrac{1+(ap_j+b+1)\mathbf{mod}3}{3}\right\rfloor\\ &= \begin{cases} 1+0=1 & \text{if} \quad (ap_j+b)\mathbf{mod}3=0\\ 0+1=1 & \text{if} \quad (ap_j+b)\mathbf{mod}3=1\\ 0+0=0 & \text{if} \quad (ap_j+b)\mathbf{mod}3=2.\end{cases}\end{aligned}$$

So $\varepsilon_1+\varepsilon_2\le 1$ and $\varepsilon'' = \left\lfloor \frac{3b}{p_j}\right\rfloor - \varepsilon_1-\varepsilon_2 \ge 0$.

□

Therefore,

$$\begin{aligned}\varepsilon &= \varepsilon' - a + \varepsilon'' + \left\lfloor \tfrac{ap_j+b+\theta_{j,i}}{p_ip_j}\right\rfloor + \left\lfloor \tfrac{ap_j+b+2}{p_ip_j}\right\rfloor\\ &\ge 1-a+\left\lfloor \tfrac{ap_j+b+\theta_{j,i}}{p_ip_j}\right\rfloor.\end{aligned}$$

If $a\le 1$ then $\varepsilon \ge 1-a\ge 0$.
Else for $a=2(<p_i)$, because $\theta_{j,i}=p_ip_j-\lambda p_j\ge p_j$ $(1\le\lambda\le p_i-1)$, $\left\lfloor \frac{ap_j+b+\theta_{j,i}}{p_ip_j}\right\rfloor \ge \left\lfloor \frac{2p_j+p_j}{3p_j}\right\rfloor = 1$. So $\varepsilon\ge 1-a+1\ge 0$.

In summary, $\varepsilon\ge 0$ for all $p_i<p_j\le\sqrt{m}$. □

Proof of (4.4). when $p_i=2$, $\left[\frac{\widetilde{1}}{p_i}\right]\equiv 0$, $m=2sp_j+t$, $t=ap_j+b$, $0\le a\le 1$, $0\le b\le p_j-1$, $s\ge 1$.

$$\begin{aligned}\varepsilon &= m\left[1-\tfrac{1}{2}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right] - \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \left[1-\tfrac{1}{2}\right]\\ &= 2sp_j\left(1-\tfrac{1}{2}\right)\left(1-\tfrac{2}{p_j}\right) - 2sp_j\left(1-\tfrac{1}{2}\right)\left(1-\tfrac{3}{p_j}\right)\\ &\quad +t-\left\lfloor\tfrac{t}{2}\right\rfloor - \left\lfloor\tfrac{t}{p_j}\right\rfloor + \left\lfloor\tfrac{t}{2p_j}\right\rfloor - \left\lfloor\tfrac{t+2}{p_j}\right\rfloor + \left\lfloor\tfrac{t+\theta_{i,j}}{2p_j}\right\rfloor\\ &\quad -\left(t-\left\lfloor\tfrac{3t}{p_j}\right\rfloor\right) + \left\lfloor\tfrac{t-\left\lfloor\frac{3t}{p_j}\right\rfloor}{2}\right\rfloor\\ &= s+\left\lfloor\tfrac{3t}{p_j}\right\rfloor - \left\lfloor\tfrac{t}{p_j}\right\rfloor - \left\lfloor\tfrac{t+2}{p_j}\right\rfloor + 0 + \left\lfloor\tfrac{t+\theta_{i,j}}{2p_j}\right\rfloor - \left\lfloor\tfrac{\left\lfloor\frac{3t}{p_j}\right\rfloor}{2}\right\rfloor - \varepsilon_1\\ &= s+3a+\left\lfloor\tfrac{3b}{p_j}\right\rfloor - a - a - \left\lfloor\tfrac{b+2}{p_j}\right\rfloor + \left\lfloor\tfrac{ap_j+b+\theta_{i,j}}{2p_j}\right\rfloor - \left\lfloor\tfrac{3a+\left\lfloor\frac{3b}{p_j}\right\rfloor}{2}\right\rfloor - \varepsilon_1\\ &= s+\left\lfloor\tfrac{3b}{p_j}\right\rfloor + \left\lfloor\tfrac{ap_j+b+\theta_{i,j}}{2p_j}\right\rfloor - \varepsilon_1-\varepsilon_3-\varepsilon_4,\end{aligned}$$

where

$$\begin{cases}\varepsilon_1 &= \left\lfloor\tfrac{\left\lfloor 3a+\frac{3b}{p_j}\right\rfloor\mathbf{mod}2+\left(ap_j+b-3a-\left\lfloor\frac{3b}{p_j}\right\rfloor\right)\mathbf{mod}2}{2}\right\rfloor\\ &= \left\lfloor\tfrac{\left\lfloor a+\frac{3b}{p_j}\right\rfloor\mathbf{mod}2+\left(ap_j+b-a-\left\lfloor\frac{3b}{p_j}\right\rfloor\right)\mathbf{mod}2}{2}\right\rfloor \le 1\\ \varepsilon_3 &= \left\lfloor\tfrac{a+\left\lfloor\frac{3b}{p_j}\right\rfloor}{2}\right\rfloor \le 1\\ \varepsilon_4 &= \left\lfloor\tfrac{b+2}{p_j}\right\rfloor \le 1,\end{cases}$$

$$\Delta\varepsilon = \varepsilon_1+\varepsilon_3+\varepsilon_4\le 3.$$

(1) If $\left\lfloor \frac{3b}{p_j} \right\rfloor = 0$, then $\varepsilon_3 = 0$, $\varepsilon_1 = \left\lfloor \frac{a\mathbf{mod}2+(ap_j+b-a)\mathbf{mod}2}{2} \right\rfloor$:
If $a = 0$, then $\varepsilon_1 = 0$, $\Delta\varepsilon = \varepsilon_4 \le 1$, $\varepsilon \ge s - \Delta\varepsilon \ge 0$;
Else for $a = 1$, then $\varepsilon_1 = \left\lfloor \frac{1+b\mathbf{mod}2}{2} \right\rfloor$. $\lambda_{i,j} = 2p_j - 2$, $\theta_{i,j} = 2p_j - \lambda_{i,j} = 2$.

$$\left\lfloor \frac{ap_j+b+\theta_{i,j}}{2p_j} \right\rfloor - \varepsilon_4 = \left\lfloor \frac{p_j+b+2}{2p_j} \right\rfloor - \left\lfloor \frac{b+2}{p_j} \right\rfloor = \begin{cases} 0-0=0 & \text{if} \quad b+2 < p_j \\ 1-1=0 & \text{if} \quad b+2 \ge p_j, \end{cases}$$

$$\left\lfloor \frac{3b}{p_j} \right\rfloor - \varepsilon_3 = \left\lfloor \frac{3b}{p_j} \right\rfloor - \left\lfloor \frac{a+\left\lfloor \frac{3b}{p_j} \right\rfloor}{2} \right\rfloor \ge \begin{cases} 0-0=0 & \text{if} \quad \left\lfloor \frac{3b}{p_j} \right\rfloor = 0 \\ 1-1=0 & \text{if} \quad \left\lfloor \frac{3b}{p_j} \right\rfloor = 1 \\ 2-1=1 & \text{if} \quad \left\lfloor \frac{3b}{p_j} \right\rfloor = 2, \end{cases}$$

So $\varepsilon = s + \left(\left\lfloor \frac{3b}{p_j} \right\rfloor - \varepsilon_3 \right) + \left(\left\lfloor \frac{ap_j+b+\theta_{i,j}}{2p_j} \right\rfloor - \varepsilon_4 \right) - \varepsilon_1 \ge 1 + 0 + 0 - 1 = 0$.

(2) If $\left\lfloor \frac{3b}{p_j} \right\rfloor = 1$, then $\varepsilon \ge s + 1 + 0 - \varepsilon_1 - \varepsilon_3 - 1 \ge 1 - \varepsilon_1 - \varepsilon_3$:
If $a = 0$, then $\varepsilon_3 = 0$, $\varepsilon \ge 1 - \varepsilon_1 - 0 \ge 0$;
Else for $a = 1$, then $\varepsilon_1 = 0$, $\varepsilon \ge 1 - 0 - \varepsilon_3 \ge 0$.

(3) If $\left\lfloor \frac{3b}{p_j} \right\rfloor = 2$, then $\varepsilon \ge s + 2 + 0 - \varepsilon_1 - \varepsilon_3 - \varepsilon_4 \ge 0$.

In summary, $\varepsilon \ge 0$ for all $2 = p_i < p_j \le \sqrt{m}$. □

Eq. (4.3) means that, after deleted the items of $Z_k\mathbf{mod}p_j = 0, p_j - 2$ from Z, the items of $Z_k\mathbf{mod}p_i \ne 0, p_i - 2$ will have at least,

$$\begin{aligned} &S(m, p_i \nparallel 0, p_i - 2; p_j \nparallel 0, p_j - 2) \\ &= S(\lceil m(1 - 3/p_j) \rceil, p_i \nparallel 0, p_i - 2) + \Delta(p_i \nparallel 0, p_i - 2), \end{aligned} \tag{4.5}$$

where $\Delta(p_i \nparallel 0, p_i - 2) \ge 0$ is the extra items of $Z_k\mathbf{mod}p_i \ne 0, p_i - 2$ and $Z_k\mathbf{mod}p_j \ne 0, p_j - 2$. Thus the effect of $\left[1 - \frac{1}{p_j} - \frac{\widetilde{1}}{p_j}\right]$ is that it constructs a new effective nature sequence with at least $\lceil m(1 - 3p_j) \rceil$ items which satisfy the condition $Z_k\mathbf{mod}p_j \ne 0, p_j - 2$.

This lemma means that, from equation (4.1) and (3.5), we can let $m\left[1 - \frac{1}{p_j} - \frac{\widetilde{1}}{p_j}\right] = \left\lceil m\left(1 - \frac{3}{p_j}\right) \right\rceil + t, t \ge 0$, and operated by $\left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right]$ to get the items which have no multiples of p_i and p_j in Z andZ'.

$$\begin{aligned} &m\left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right]\left[1 - \frac{1}{p_j} - \frac{\widetilde{1}}{p_j}\right] = S(m, p_i \nparallel 0, p_i - 2; p_j \nparallel 0, p_j - 2) \\ &= \left(\left\lceil m\left(1 - \frac{3}{p_j}\right)\right\rceil + t\right)\left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right] \\ &= S(\lceil m(1 - 3/p_j) \rceil, p_i \nparallel 0, p_i - 2) + \Delta(p_i \nparallel 0, p_i - 2) \\ &= \left\lceil m\left(1 - \frac{3}{p_j}\right)\right\rceil\left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right] + t\left[1 - \frac{1'}{p_i} - \frac{1''}{p_i}\right] \\ &\ge \left\lceil m\left(1 - \frac{3}{p_j}\right)\right\rceil\left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i}\right]. \end{aligned} \tag{4.6}$$

Lemma 4.4.

$$m\left[1 - \tfrac{1}{2}\right]\left[1 - \tfrac{1}{3} - \tfrac{\widetilde{1}}{3}\right] \ge \left\lceil \tfrac{m}{6} \right\rceil - 1. \tag{4.7}$$

Proof. For $p_i = 2, p_j = 3$, $\frac{\widetilde{1}}{2} = 0$, let $m = 6s + t$,

$$\begin{aligned} &m\left[1 - \tfrac{1}{2}\right]\left[1 - \tfrac{1}{3} - \tfrac{\widetilde{1}}{3}\right] = 6s\left[1 - \tfrac{1}{2}\right]\left[1 - \tfrac{1}{3} - \tfrac{\widetilde{1}}{3}\right] + t\left[1 - \tfrac{1}{2}\right]\left[1 - \tfrac{1}{3} - \tfrac{\widetilde{1}}{3}\right] \\ &\ge 6s\left(1 - \tfrac{1}{2}\right)\left(1 - \tfrac{2}{3}\right) = s = \left\lfloor \tfrac{m}{6} \right\rfloor \ge \left\lceil \tfrac{m}{6} \right\rceil - 1. \end{aligned}$$

For the residual class of modulo 6, $X = \{6s+1, 6s+2, \cdots, 6s+6\}$, $X_k\mathbf{mod}6 = \{1,2,3,4,5,6\}$, there are 4 elements $(2,3,4,6)$ of multiples of 2 or 3. For the other elements $X_k\mathbf{mod}6 = \{1,5\}$, there is at least one with $X_k\mathbf{mod}6 = 5$, i.e., $X_k\mathbf{mod}2 \neq 0, X_k\mathbf{mod}3 \neq 0$, and $X_k\mathbf{mod}3 \neq (3-2) = 1$. The twin primes have at least $\lfloor \frac{m}{6} \rfloor \geq \lceil \frac{m}{6} \rceil - 1$, or $m\left[1-\frac{1}{2}\right]\left[1-\frac{1}{3}-\frac{\widetilde{1}}{3}\right] \geq \lceil \frac{m}{6} \rceil - 1$. □

Lemma 4.5. *For* $i = 1, 2, \cdots, i_m, j$,

$$m\prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right] \geq \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right], \tag{4.8}$$

where $\frac{\widetilde{1}}{p_i} \equiv 0$ *for* $p_i = 2$.

Proof. From equation (2.16),(4.1),(4.3)-(4.6), For $i_1 \neq i_2$, we have $S(m, p_{i_1} \nparallel 0, p_{i_1}-2; p_j \nparallel 0, p_j-2) \geq S(\lceil m(1-3/p_j)\rceil, p_{i_1} \nparallel 0, p_{i_1}-2)$, $S(m, p_{i_2} \nparallel 0, p_{i_2}-2; p_j \nparallel 0, p_j-2) \geq S(\lceil m(1-3/p_j)\rceil, p_{i_2} \nparallel 0, p_{i_2}-2)$, so that $S(m, p_{i_1} \nparallel 0, p_{i_1}-2; p_{i_2} \nparallel 0, p_{i_2}-2; p_j \nparallel 0, p_j-2) \geq S(\lceil m(1-3/p_j)\rceil, p_{i_1} \nparallel 0, p_{i_1}-2; p_{i_2} \nparallel 0, p_{i_2}-2)$. $\cdots\cdots$. $S(m, p_{i_1} \nparallel 0, p_{i_1}-2; p_{i_2} \nparallel 0, p_{i_2}-2; \cdots, p_{i_m} \nparallel 0, p_{i_m}-2; p_j \nparallel 0, p_j-2) \geq S(\lceil m(1-3/p_j)\rceil, p_{i_1} \nparallel 0, p_{i_1}-2; p_{i_2} \nparallel 0, p_{i_2}-2; \cdots, p_{i_m} \nparallel 0, p_{i_m}-2)$.

$$\begin{aligned}
&m\prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]\\
&= S(m, p_1 \nparallel 0; p_2 \nparallel 0, p_2-2; \cdots; p_{i_m} \nparallel 0, p_{i_m}-2; p_j \nparallel 0, p_j-2)\\
&\geq S(\lceil m(1-3/p_j)\rceil, p_1 \nparallel 0; p_2 \nparallel 0, p_2-2; \cdots; p_{i_m} \nparallel 0, p_{i_m}-2)\\
&= \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right].
\end{aligned} \tag{4.9}$$

Besides, suppose that for $1 < r \leq i_m$,

$$m\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right] = \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] + t, \tag{4.10}$$

where $t \geq 0$. It means that the effect of the operator $\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]$ when operating on m is that dividing $Z = \{1, 2, \cdots, m\}$ into two set $X = \{1, 2, \cdots, m' = \left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\}$ and $X' = \{X'_1, X'_2, \cdots, X'_t, t = m - m' \geq 0\}$. From equation (2.18), (2.19), (4.1), (4.3), and (4.10), we have

$$\begin{aligned}
&m\prod_{i=r-1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]\\
&= m\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]\left[1-\tfrac{1}{p_{r-1}}-\tfrac{\widetilde{1}}{p_{r-1}}\right]\\
&= \left(\left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] + t\right)\left[1-\tfrac{1'}{p_{r-1}}-\tfrac{1''}{p_{r-1}}\right]\\
&\geq \left(\left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\right)\left[1-\tfrac{1'}{p_{r-1}}-\tfrac{1''}{p_{r-1}}\right]\\
&= \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \prod_{i=r-1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right].
\end{aligned}$$

Or

$$\begin{aligned}
&m\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]\left[1-\frac{1}{p_{r-1}}-\frac{\widetilde{1}}{p_{r-1}}\right]\\
&=m\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]-m\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]\left[\frac{1}{p_{r-1}}\right]\\
&\quad-m\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]\left[\frac{\widetilde{1}}{p_{r-1}}\right]\\
&=\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]+t\\
&\quad-\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[\frac{1}{p_{r-1}}\right]-t\left[\frac{1'}{p_{r-1}}\right]\\
&\quad-\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[\frac{\widetilde{1}}{p_{r-1}}\right]-t\left[\frac{1''}{p_{r-1}}\right]\\
&=\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_{r-1}}-\frac{\widetilde{1}}{p_{r-1}}\right]+t\left[1-\frac{1'}{p_{r-1}}-\frac{1''}{p_{r-1}}\right]\\
&\ge\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\prod_{i=r-1}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right].
\end{aligned}$$

In fact, if

$$m\prod_{i=1}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]<\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\prod_{i=1}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right],$$

then for any p_i, p_j, before deleting the multiples of other primes, it must have

$$m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]<\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right].$$

which contradicts Lemma 4.3. So this lemma is true.

If $p_{r-1}=2$, then

$$\begin{aligned}
&m\left[1-\frac{1}{p_{r-1}}\right]\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]\\
&\ge\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\prod_j\left[1-\frac{1}{p_{r-1}}\right]\prod_{i=r}^{i_m}\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right].
\end{aligned}\tag{4.11}$$

$\square$

With Lemma 4.3, the operator $\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]$ can be represented by $\left(1-\frac{3}{p_j}\right)$, and other operator $\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]$ can operate on this inequality unchanged.

5. EXPLANTATION

Let $m=sp_ip_j+ap_j+b\ge p_j^2$, then for all $p_i<p_j$, $m\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]$ is effective to a nature sequence $X=\{f(Z)\}$ whose number is not less than $\lceil m(1-3/p_j)\rceil$. The reason is as follows. When a nature sequence is deleted by the items of $Z_k\mathbf{mod}p_j=0, p_j-2$, the sequence is subtracted by $\left[\frac{m}{p_j}\right]+\left[\frac{m+2}{p_j}\right]$. We can arrange the m items in a table of p_j rows (Table 1). $\left[\frac{m}{p_j}\right]$ will delete the p_jth row, and $\left[\frac{m+2}{p_j}\right]$ will delete the $(p_j-2)th$ row. Thus there are (p_j-2) rows left in which each item $Z_k\mathbf{mod}p_j\neq 0, p_j-2$.

Table 1. Set Z

1	p_j+1	$\cdots$	$(p_i-1)p_j+1$	$\cdots$	$(sp_i-1)p_j+1$	sp_ip_j+1	$\cdots$	$sp_ip_j+ap_j+1$
2	p_j+2	$\cdots$	$(p_i-1)p_j+2$	$\cdots$	$(sp_i-1)p_j+2$	sp_ip_j+2	$\cdots$	$sp_ip_j+ap_j+2$
$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$
b	p_j+b	$\cdots$	$(p_i-1)p_j+b$	$\cdots$	$(sp_i-1)p_j+b$	sp_ip_j+b	$\cdots$	$sp_ip_j+ap_j+b$
$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	
p_j	$2p_j$	$\cdots$	p_ip_j	$\cdots$	sp_ip_j	$sp_ip_j+p_j$	$\cdots$	

But every p_i items ($0 \le Z_k\mathbf{mod}p_i \le p_i-1$) in any row of the first sp_i columns consist in a complete system of residues modulo p_i, because $C_1 = \{1, p_j+1, 2p_j+1, \cdots, (p_i-1)p_j+1\}$ and $C_r = \{C_1 + r\}$ are both complete system of residues modulo p_i, where r is any (row or column) constant. There are (p_j-2) such rows or $sp_i(p_j-2)$ items left. These items are effective to a nature sequence when deleting the items of $Z_k\mathbf{mod}p_i = 0, p_i-2$.

$sp_ip_j\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right] = sp_i(p_j-2)\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right] = s(p_i-2)(p_j-2).$

Let $t = ap_j+b$, $0 \le b \le p_j-1$, $0 \le a \le p_i-1$. $t\left[\frac{1}{p_j}+\frac{\widetilde{1}}{p_j}\right]$ will delete at most $a+(a+1) \le t\frac{1}{p_j} + sp_ip_j\frac{1}{p_j} = \frac{m}{p_j}$ items. If we add these items by removing those from the end of sequence then the sequence is again effective to a nature sequence, the sequence left has at least,

$$\begin{aligned}
M(j) &\ge sp_i(p_j-2) + t\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right] - [a+(a+1)] \\
&\ge sp_i(p_j-3) + t + sp_i - 4a - 2 \\
&= sp_i(p_j-3) - 3\left\lfloor \frac{t}{p_j} \right\rfloor + t + (sp_i - a - 2) \\
&\ge sp_ip_j(1-\frac{3}{p_j}) + \left\lceil t - \frac{3t}{p_j} \right\rceil + (sp_i - a - 2) \\
&= \left\lceil (sp_ip_j + t)(1-\frac{3}{p_j}) \right\rceil + (sp_i - a - 2) \\
&= \left\lceil m(1-\frac{3}{p_j}) \right\rceil + (sp_i - a - 2).
\end{aligned}$$

For $s \ge 2$ or $a \le p_i - 2$, we have $(sp_i - a - 2) = (s-1)p_i + (p_i - a - 1) - 1 \ge 0$, $M(j) \ge \left\lceil m(1-\frac{3}{p_j}) \right\rceil$.

For $s = 1$ and $a = p_i - 1$, the items of t have p_j rows, $p_i - 1$ columns and some b items. In each of the first b rows, there are exact p_i items which consist in a complete system of residues modulo p_i, and these items can be considered as an effective nature sequence when deleting the multiples of p_i ($Z_k\mathbf{mod}p_i = 0, p_i-2$). The other items have at most p_j rows and $p_i - 1$ columns where the multiples of p_j have at most $2(p_i-1)$. As before, we can add these items to make the t as an effective nature sequence, therefore,

$$\begin{aligned}
M(j) &\ge sp_i(p_j-2) + t\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right] - 2(p_i-1) \\
&\ge \left\lceil m(1-\frac{3}{p_j}) \right\rceil + (sp_i - a - 1) \ge \left\lceil m(1-\frac{3}{p_j}) \right\rceil.
\end{aligned}$$

Thus for any $p_i < p_j$, the original sequence of $m \geq p_j^2$, when deleted by the items of $Z_k \mathbf{mod} p_j = 0, p_j - 2$ from Z, is effective to reconstruct a new natural sequence having at least $\left\lceil m(1 - \frac{3}{p_j}) \right\rceil$ items.

Example 5.1. *$n = 41$, $P = \{2, 3, 5\}$, $Z = \{1, 2, \cdots, 41\}$.*

For $p_j = 5$, after deleted the item of $Z_k \mathbf{mod} p_j = 0, 5 - 2$, it becomes $Z \to Z' = \{1, 2, 4, 6, 7, 9, 11, 12, 14, 16, 17, 19, 21, 22, 24, 26, 27, 29, 31, 32, 34, 36, 37,\text{-}39, 41\}$. we can rearrange these items as $Z' = \{1, 2, (21), 4, (29), 6, 7, (26), 9, (22),\text{-}11, 12, (19), 14, (27), 16, 17||, 24, 31, 32, 34, 36, 37, 39, 41\}$. The first $\left\lceil n\left(1 - \frac{3}{p_j}\right) \right\rceil = 17$ items can be taken as an effective natural sequence from the original one when deleting the items of $Z_k \mathbf{mod} 3 = 0, 3 - 2$. The other sequence $X = \{24, 31, 32, 34, 36,\text{-}37, 39, 41\}$, having at least zero item when deleting the multiples of all primes, will be neglected in further procession.

6. Proof of Theorem 1.1

For a given n, consider the possible twin primes of $[Z_k, Z_k + 2]$ in set $Z = \{1, 2, \cdots, n\}$, $p_v^2 \leq n < p_{v+1}^2$.

Lemma 6.1. *The number of twin prime pairs in n,*

$$D(n) \geq \left\lceil \frac{p_v}{3} \prod_{i=3}^{v-1} \frac{p_{i+1} - 3}{p_i} \right\rceil + D(\sqrt{n}) - 2. \tag{6.1}$$

Proof. Because $m(v) = n \geq p_v^2$. Let $m(j-1) = \lceil m(j)(1 - 3/p_j) \rceil$, then

$$\begin{aligned} m(j-1) &= \lceil m(j)(1 - 3/p_j) \rceil \geq p_j^2(1 - 3/p_j) = p_j(p_j - 3) \\ &\geq (p_{j-1} + 2)(p_{j-1} - 1) = p_{j-1}^2 + p_i - 2 \geq p_{j-1}^2. \end{aligned}$$

And for any $i \leq (j-1)$, we have $m(i) \geq p_i^2$.

From equation (2.12), (4.8), (4.11) and (4.7),

$$\begin{aligned} D_0(n) &= n \left[1 - \tfrac{1}{2}\right] \prod_{i=2}^{v} \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i}\right] \\ &\geq \left\lceil n\left(1 - \tfrac{3}{p_v}\right) \right\rceil \left[1 - \tfrac{1}{2}\right] \prod_{i=2}^{v-1} \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i}\right] \\ &\geq \left\lceil \left\lceil n\left(1 - \tfrac{3}{p_v}\right) \right\rceil \left(1 - \tfrac{3}{p_{v-1}}\right) \right\rceil \left[1 - \tfrac{1}{2}\right] \prod_{i=2}^{v-2} \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i}\right] \\ &\geq \cdots \\ &\geq \left\lceil n \prod_{i=3}^{v} \left(1 - \tfrac{3}{p_i}\right) \right\rceil \left[1 - \tfrac{1}{2}\right] \left[1 - \tfrac{1}{p_2} - \tfrac{\tilde{1}}{p_2}\right] \\ &\geq \left\lceil \tfrac{n}{6} \prod_{i=3}^{v} \left(1 - \tfrac{3}{p_i}\right) \right\rceil - 1 \end{aligned}$$

but

$$\begin{aligned} \prod_{i=3}^{v} \left(1 - \tfrac{3}{p_i}\right) &= \tfrac{p_3 - 3}{p_3} \tfrac{p_4 - 3}{p_4} \cdots \tfrac{p_{v-1} - 3}{p_{v-1}} \tfrac{p_v - 3}{p_v} \\ &= \tfrac{p_3 - 3}{p_v} \tfrac{p_4 - 3}{p_3} \tfrac{p_5 - 3}{p_4} \cdots \tfrac{p_v - 3}{p_{v-1}} = \tfrac{2}{p_v} \prod_{i=3}^{v-1} \tfrac{p_{i+1} - 3}{p_i}, \end{aligned}$$

and $n \geq p_v^2$, so

$$D_0(n) \geq \left\lceil \frac{n}{3p_v} \prod_{i=3}^{v} \left(1 - \frac{3}{p_i}\right) \right\rceil - 1 \geq \left\lceil \frac{p_v}{3} \prod_{i=3}^{v-1} \frac{p_{i+1} - 3}{p_i} \right\rceil - 1.$$

From equation (2.13),(2.14),

$$\begin{aligned} D(n) &= D_0(n) + D(\sqrt{n}) - D_1 \\ &\geq \left\lceil \tfrac{p_v}{3} \prod_{i=3}^{v-1} \tfrac{p_{i+1}-3}{p_i} \right\rceil - 1 + D(\sqrt{n}) - D_1 \\ &\geq \left\lceil \tfrac{p_v}{3} \prod_{i=3}^{v-1} \tfrac{p_{i+1}-3}{p_i} \right\rceil + D(\sqrt{n}) - 2. \end{aligned}$$

□

Proof of Theorem 1.1. Suppose that there is no twin prime when greater than enough large number n_M.

$$D(n) \quad \leq D(n_M) \quad \text{for enough } n > n_M. \tag{6.2}$$

Consider the pairs of twin prime in the range $[1, n]$, where $n = n_M^2$,

$$D(n) \quad \geq \left\lceil \tfrac{p_V}{3} \prod_{i=3}^{V-1} \tfrac{p_{i+1}-3}{p_i} \right\rceil + D(\sqrt{n}) - 2. \tag{6.3}$$

where p_V is the maximum prime in n.

Then the twin prime between n_M and $n = n_M^2$ will have,

$$\begin{aligned} \Delta D(n) &= D(n) - D(n_M) = D(n) - D(\sqrt{n}) \\ &\geq \left\lceil \tfrac{p_V}{3} \prod_{i=3}^{V-1} \tfrac{p_{i+1}-3}{p_i} \right\rceil - 2 = \left\lceil \tfrac{W(V)}{3} \right\rceil - 2. \end{aligned} \tag{6.4}$$

where

$$W(V) = p_V \prod_{i=3}^{V-1} \frac{p_{i+1} - 3}{p_i}. \tag{6.5}$$

If $W(V) > 6$ then $\Delta D(n) \geq 1$, there will be at least one twin prime between n_M and $n = n_M^2$.

For $V \geq V_0 = 5, p_{V_0} = 11$,

$$W(5) = 11 \frac{4}{5} \frac{8}{7} = 10.057 > 6 \tag{6.6}$$

Suppose that for V, $W(V) = p_V \prod_{i=3}^{V-1} \frac{p_{i+1}-3}{p_i} > 6$, then for $V + 1$,

$$\begin{aligned} W(V+1) &= p_{V+1} \prod_{i=3}^{V-1} \tfrac{p_{i+1}-3}{p_i} \tfrac{p_{V+1}-3}{p_V} \\ &= p_V \prod_{i=3}^{V-1} \tfrac{p_{i+1}-3}{p_i} \tfrac{p_{V+1}}{p_V} \tfrac{p_{V+1}-3}{p_V} \\ &= W(V) \tfrac{p_{V+1}^2 - 3p_{V+1}}{p_V^2}. \end{aligned} \tag{6.7}$$

Because, $p_{V+1} = p_V + 2\Delta, \Delta \geq 1, p_V \geq 5$,

$$\begin{aligned} & p_{V+1}^2 - 3p_{V+1} - p_V^2 = (p_V + 2\Delta)^2 - 3(p_V + 2\Delta) - p_V^2 \\ & = p_V^2 + 4\Delta p_V + 4\Delta^2 - 3p_V - 6\Delta - p_V^2 \\ & = 3(\Delta - 1)p_V + \Delta(p_V + 4\Delta - 6) > 0. \end{aligned}$$

Therefore, $\frac{p_{V+1}^2 - 3p_{V+1}}{p_V^2} > 1$, and

$$W(V+1) > W(V) > 6. \tag{6.8}$$

So that $\Delta D(n) \geq 1$. It contracts the supposition of Eq. (6.2). Therefore ‘there are infinitely many pairs of twin prime’. From Eq. (6.4) and (6.5), $\Delta D(n)$ approaches infinity as n grows without bound. The proof is completed. □

Example 6.2 (Actual vs. Simplified Formula)**.** Let

$$D'(n) \quad = \left\lceil \tfrac{p_v}{3} \prod_{i=3}^{v-1} \tfrac{p_{i+1}-3}{p_i} \right\rceil - 2. \tag{6.9}$$

Then from Eq. (6.3), it should have $D(n) \geq D'(n)$.

Figure 1 shows the actual pairs of twin prime $D(n)$ (solid line) in the range of $[1, p_v^2 + 1]$, and its simplified formula $D'(n)$(dashed line) from Eq. (6.9). It clearly shows that $D(n) \geq D'(n)$, and $D'(n)$ has no up bound for enough large n.



Figure 1. The minimum number of actual prime pairs (solid) and its simplified formula (dashed) against v.



Figure 2. The ratio of $r = D(n)/D'(n)$ against v.

Figure 2 shows the ratio of

$$r = \tfrac{D(n)}{D'(n)}. \tag{6.10}$$

It shows that, this ratio is greater than one and seems to progress as n increase. It also shows that the formula (Eq. 6.9) used in our proof is only a small part of the actual twin prime pairs.

References

[1] Kenneth H. Rosen, *Elementanry Number Theory and Its Applications*, Fifth Edition, Pearson Education Asia Limited and China Maxhine Press, 2005.
[2] Hardy, G. H. and Littlewood, J. E. *Some Problems of 'Partitio Numerorum.* III. On the Expression of a Number as a Sum of Primes." Acta Math. **44**(1923), 1-70.
[3] K. Indlekofer and A. Jrai, *Largest known twin primes,* Math. Comp., **65** (1996) 427-428.
[4] M. Suzuki, *Alternative formulations of the twin prime problem,* Amer. Math. Monthly, **107:1** (2000) 55–56.

Chinese Academy of Sciences, A20 Datun rd. Chaoyang Dst. Beijing 100012, China
E-mail address: `zldu@bao.ac.cn`

Chinese Academy of Sciences, 99 Donggang rd. ShiJiaZhuang, HeBei, 050031, China
E-mail address: `shouyudu@yahoo.com.cn`